

NEMATICIDE

FOSTHIAZATE

Contact nematicide

Fosthiazate is a nematicide discovered and developed by ISK.

Fosthiazate provides a good and stable control of cyst, root-knot, root lesion and free-living nematodes in a wide range of crops such as potatoes, bananas, tomatoes, and other vegetables.


The control activities of fosthiazate remain unaffected by soil type, soil pH and soil temperature.

Since the early 1990's, ISK has commercialized fosthiazate in multiple countries.



Physico-Chemical Properties

Chemical structure



Class : organophosphate

IUPAC name : (RS)-S-sec-butyl-O-ethyl-2-oxo-1,3-thiazolidin-3-ylphosphonothioate

Molecular weight : 283.3

Molecular formula : C₉H₁₉NO₃PS₂

Vapour pressure : 5.6 × 10⁻¹ mPa (25°C)

Water solubility : 9.00 g/L (pH 7, 25°C)

Form : Liquid, slightly yellow

Development code : IKI-1145

Toxicology & Ecotoxicology

Rat LD₅₀ (oral) : 73 mg/kg (m), 57 mg/kg (f)

Rat LD₅₀ (dermal) : 2,396 mg/kg (m), 861 mg/kg (f)

Rat LC₅₀ (inhalation) : 0.83 mg/L (m), 0.56 mg/L (f)

Skin irritation : slight irritant (rabbit)

Eye irritation : irritant (rabbit)

Skin sensitization : sensitizer (guinea pig)

Avian LD₅₀ (acute oral) : 10 mg/kg bw (quail)

Avian LD₅₀ (acute oral) : 20 mg/kg (mallard duck)

Fish LC₅₀ : 114 mg/L (trout, 96 h)

Fish LC₅₀ : 171 mg/L (bluegill, 96 h)

Bees LD₅₀ (acute oral) : 0.61 µg/bee (48 h)

Bees LD₅₀ (acute contact) : 0.26 µg/bee (48 h)

Daphnia magna EC₅₀ : 0.47 mg/L (48 h)

Application

Broadcast and soil incorporation to control nematodes (*Meloidogyne* spp., *Heterodera* spp. and *Pratylenchus* spp.), aphids, mites, thrips, etc in vegetables, potatoes, tomatoes and bananas, etc. Applied mainly at 1.5-3.0 kg a.i./ha.

Mode of Action

Inhibition of acetylcholine esterase in nematodes and other pests. Its excellent systemic action also provides high performance against nematodes and foliar insect pests. Acts against motile larval stages of nematodes in the soil and prevents invasion to roots of crops.

Product

Trade Names	NEMATHORIN, CIERTO, NEMABUSTER, 福氣多, etc.	
Formulations	1.5%G, 5%G, 10%G, 30%SL, 150EC	
Registered Countries	Asia	China, Japan, South Korea, Taiwan, Philippines, etc.
	Europe	Belgium, France, Germany, Hungary Romania, Italy, Morocco, Netherlands, Spain, Turkey, UK, etc.
Crops	Vegetables, Potatoes, Sweet potatoes, Bananas, Pine tree, etc.	



ISHIHARA SANGYO KAISHA, LTD.

URL : <http://www.iskweb.co.jp> E-mail : isk.bio@iskweb.co.jp
1-3-15 Edobori, Nishi-ku, Osaka 550-0002 TEL +81-6-6444-7154

The Efficacy of Fosthiazate on Root-knot nematodes in Tomato



Untreated



Fosthiazate (2 kg a.i./ha)

Damaged Melon by *Meloidogyne incognita*



Untreated

Fosthiazate

Characteristics

Excellent control of major plant parasitic nematode pests such as root-knot, root-lesion, cyst, burrowing nematodes

It exhibits nematocidal and nemastatic activities.

Stable activities in various soil conditions (soil pH, temperature, moisture, etc.)

Improves yield and quality of crops

Excellent selectivity on crops

Applied at relatively lower dose rates compared with other soil applied insecticides & nematicide (1.5 to 3.0 kg a.i./ha)

Pest Spectrum

Nematodes	Root-knot nematode	<i>Meloidogyne</i> spp.
	Root-lesion nematode	<i>Pratylenchus</i> spp.
	Cyst nematode	<i>Globodera</i> spp., <i>Heterodera</i> spp.
	Burrowing nematode	<i>Radopholus similis</i>
	Potato rot nematode, Iris nematode	<i>Ditylenchus destructor</i>
	Spiral nematode	<i>Helicotylenchus</i> sp.
	Stylet nematode	<i>Tylenchorhynchus</i> spp.
	Chrysanthemum folier nematode	<i>Aphelenchoides ritzemabosi</i>
	Pine wood nematode	<i>Bursaphelenchus xylophilus</i>
	Rice white-tip nematode	<i>Aphelenchoides besseyi</i> Christie
Other insects	Thrips, Aphids, Mites, Whiteflies, Bugs, Beetles	



Meloidogyne incognita



Globodera rostochiensis

